



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ON PROBLEM 132.

BY PROF. A. HALL, WASHINGTON, D. C.

THE integral required in this question is a special case of the one discussed by Legendre in his treatise on elliptic functions, Chap. XXVI. p. 165. Legendre shows that the elliptic integrals disappear, and that the result can be expressed in logarithmic and circular functions. His method is complicated however, and Clausen has obtained the result by the following simple substitutions. Let

$$z = \frac{x-1}{\sqrt{(x^3-1)}}: \quad z' = \sqrt{(x^3-1)}: \quad z'' = \frac{(x-1)^2}{\sqrt{(x^3-1)}}:$$

Then

$$dz = -\frac{x^3-3x^2+2}{2\sqrt{(x^3-1)^3}}dx: \quad dz' = \frac{3}{2} \cdot \frac{x^2 \cdot dx}{\sqrt{(x^3-1)}}:$$

$$dz'' = \frac{1}{2} \cdot \frac{x^4+2x^3-3x^2-4x+4}{\sqrt{(x^3-1)^3}}dx:$$

Hence

$$\frac{dz}{1-3z^2} = -\frac{1}{2} \cdot \frac{x^2-2x-2}{x^2-2x+4} \cdot \frac{dx}{\sqrt{(x^3-1)}}: \quad \frac{dz'}{z'^2+9} = \frac{3}{2} \cdot \frac{x^2}{x^3+8} \cdot \frac{dx}{\sqrt{(x^3-1)}}:$$

$$\frac{dz''}{z''^2+9} = \frac{1}{2} \cdot \frac{(x-1)(x^3+3x^2-4)}{(x+2)(x^3+3x^2-4)} \cdot \frac{dx}{\sqrt{(x^3-1)}} = \frac{1}{2} \cdot \frac{x-1}{x+2} \cdot \frac{dx}{\sqrt{(x^3-1)}}:$$

$$\therefore \quad \frac{dz}{1-3z^2} + \frac{dz'}{z'^2+9} + \frac{dz''}{z''^2+9} = \frac{bx \cdot dx}{(x^3+8)\sqrt{(x^3-1)}}:$$

The integrations can now be performed and we have

$$\begin{aligned} \int \frac{x dx}{(x^3+8)\sqrt{(x^3-1)}} &= \frac{1}{12\sqrt{3}} \cdot \log \left(\frac{1+z\sqrt{3}}{1-z\sqrt{3}} \right) + \frac{1}{18} \tan^{-1} \frac{z'}{3} + \frac{1}{18} \tan^{-1} \frac{z''}{3} \\ &= \frac{1}{12\sqrt{3}} \cdot \log \frac{\sqrt{(x^2+x+1)} + \sqrt{(x-1)} \cdot \sqrt{3}}{\sqrt{(x^2+x+1)} - \sqrt{(x-1)} \cdot \sqrt{3}} + \frac{1}{18} \tan^{-1} \frac{3x(x-1)}{(4-x)(x^3-1)^{\frac{1}{2}}}. \end{aligned}$$

REMARKS BY PROF. ORSON PRATT, SEN., SALT LAKE CITY, UTAH.
—THE six propositions, published in the ANALYST, Vol. III, No. 6, and demonstrated in the last No., are not “presented as the basis of a theory of gravitation.” They have no bearing on or reference to the *cause* of the force called gravity, or to the exploded hypothesis of an ethereal pressure, exerted in the direction of a gravitating center.